

Carlos Kusano Bucalen Ferrari^{1,2*},
Grazielle Souza Lira Ferrari², Lohany
Dourado Nery², Daniela Franco dos
Santos², Noeli Silva Pereira², Jéssica
Silva Gonçalves² and Geane Francis
Santana²

¹Curso of Public Health, Federal University of Latin
American Integration, (UNILA), Brazil

²ICBS, The Federal University of Mato Grosso,
University Campus do Araguaia. Av. Valdon
Varjão, 6390, Industrial District, 78600-000, Barra do
Garças, MT, Brazil

Dates: Received: 25 April, 2016; Accepted: 29
June, 2016; Published: 30 June, 2016

*Corresponding author: Carlos K B Ferrari,
Professor, ICBS, the Federal University of Mato
Grosso, University Campus do Araguaia. Av. Valdon
Varjão, 6390, Industrial District, 78600-000, Barra do
Garças, MT, Brazil, Tel: No: 55 45 3522-8091; 55 45
9992-6495; E-mail: ferrariphd@yahoo.com.br

www.peertechz.com

Keywords: Loneliness; Elderly; Affective bond;
Marital status

Research Article

Affective Bond, Loneliness and Socioeconomic Aspects of an Elderly Population in Midwest, Brazil

Abstract

Background: loneliness, a common feature among elderly people, can affect health and quality of life.

Objective: this article examined affective bond and marital status among elders in “Barra do Garças”, municipality, Mato Grosso State, Midwest Brazil.

Methodology: from the 3,593 elderly people, 300 engaged in the study which had been approved by the ethical committee on research.

Results: there were more illiterate men than women (37.7% and 21.6%, respectively). In respect of income source there was a relative contingent of women with no income comparing to the men. Considering the socioeconomic class the majority of elderly, 68% belong to the less fortunate socioeconomic classes (D and E).

Conclusion: it has been observed a higher contingent of elderly people with no affective bond or living alone which influence the quality of life of those elderly by increasing the psychosocial as well as the health-related risks, which was widely discussed according to the world’s scientific literature.

Introduction

Elderly population in Latin American and Caribe grows 3.8% *per annum*; In 2005, this population group comprised 34,2 million of people which will reach 100 million in 2025 [1]. Actually, the Brazilian Elderly population comprises 8.2%, representing 16,812.608 of people [2]. Rio de Janeiro, Rio Grande do Sul, São Paulo and Paraíba are Brazilian States with highest proportion of elderly, while in Mato Grosso state only 5.9% of people are elderly [2].

Due to the potential increase of the elderly population it is really necessary to know the different social, cultural, economic, and environmental aspects related to aging, since these factors can be determinants of both the health status and the functional capacity of elderly [3].

It is well known that education level of elderly people is inversely associated with risk of chronic non-communicable diseases such as dementia, depression, type 2 diabetes mellitus, obesity, hypertension, cardiovascular diseases, poor mobility, among others [4,5].

Loneliness and lacking of affective bonds, common problems of elderly men and women, are associated with increased risk of chronic diseases such as mental disorders and cardio metabolic diseases [6].

Since loneliness has been associated with risk of depression, fatigue and pain [7], the objective of the current work was evaluate affective bond and marital status among an elderly population of a municipality from Midwest Brazil.

Methodology

The current study was done in Barra do Garças municipality, Mato Grosso State, located at the border of Goiás state. The Barra do Garças

city presents 9,141,84 Km² of area and is 510Km far from Cuiabá, the state’s capital. The city population is 58,398 inhabitants and 3,593 people have 65 years old or more [2]. The studied population was constituted by people with 65 years old or more which attended at leisure-time activities offered by Social Assistance Secretary (SAS) of Barra do Garças municipality in partnership with the Federal University of Mato Grosso (UFMT).

The convenience sample (n=300) included all subjects with 65 years old and more which engaged in leisure-time activities and agreed to be interviewed. Information regarding marital status, and socioeconomic variables such as gender, age, income and other information was collected.

This transversal study was carried out during May and September 2015.

People with 65 years old, engaged in the social activities, and that agree to engage into the study were included. The exclusion criteria was being less than 65 years old and those who did not agree to participate.

This study did not present any kind of risk for the participants and the interview ensured the privacy of each subject. All participants signed the informed consent and the study was approved by the Ethics Committee on Research of the Julio Müller University Hospital (UFMT) (665/CEP-HUJM/09).

Data was statistically analyzed by the EPI-INFO[®] software (6.04d). The hypotheses of association between variables was accepted when the found *p* value was lower than 0.05.



Results

As observed in Table 1, there is a trend of femininity among the elderly population, since almost 65% of them was women.

Considering the education level, the frequency of illiterate men was higher than that found for women.

On contrary, regarding income source there was a relative number of women who did not have income compared to the men. Regarding to the socioeconomic classification there was statistically significant differences regarding gender considering belong to the B1 and B2 socioeconomic classes. It is important to note that the majority of elderly was poor and belong to the D social class (60.7%), the second lowest.

Moreover, about 30% of the elderly, regardless of gender, declared live alone (Table 1).

Regarding marital status, the frequency of untethered elderly was 50% lower than the frequencies of the elderly with bond or widowed (Table 2). The frequency of elderly without affective bond or widowed was 20.3% and 40.3%, respectively (Table 2).

Table 1: Socioeconomic profile of elderly from Barra do Garças, MT, Brazil.

| Variables | Female (n=194) | Male (n=106) | Total (n=300) | p* |
|--------------------------------|----------------|--------------|---------------|---------|
| Age (years): | | | | |
| 65 a 69 | 111 (57.2%) | 43(40.5%)* | 154 (51.3%) | p<0,05* |
| 70 a 79 | 71 (36.6%) | 49(46.2%)* | 120 (40.0%) | |
| 80 e + | 12 (6.2%) | 14 (13.2%) | 26 (8.7%) | |
| Education level (years) | | | | |
| Never | 42 (21.6%) | 40 (37.7%)* | 82 (27.3%) | p<0,05* |
| 1 a 4 | 93 (47.9%) | 50 (47.2%) | 143 (47.7%) | |
| 5 a 8 | 40 (20.6%) | 11 (10.4%) | 51 (17.0%) | |
| 8 e + | 19 (9.8%) | 5 (4.7%) | 24(8.0%) | |
| Incomed source | | | | |
| Without | 26 (13.4%) | 1 (1.0%)* | 27 (9.0%) | p<0,05* |
| Working | 16 (8.2%) | 23 (21.6%) | 39 (13.0%) | |
| Retired | 146 (75.3%) | 81 (76.4%) | 227 (75.7%) | |
| Others | 6 (3.1%) | 1 (1.0%) | 7 (2.3%) | |
| Socioeconomic classes | | | | |
| B1 and B2 | 13 (6.7%) | 3 (2.8%)* | 16 (5.3%) | p<0,05* |
| C | 53 (27.3%) | 27 (25.5%) | 80 (26.7%) | |
| D | 120 (61.9%) | 62 (58.5%) | 182 (60.7%) | |
| E | 8 (4.1%) | 14 (13.2%) | 22 (7.3%) | |
| Live alone? | | | | |
| Yes | 54 (27.8%) | 34 (32.1%) | 88 (29.3%) | p>0,05 |
| No | 86 (72.2%) | 54 (27.8%) | 212 (70.7%) | p<0,05 |

*p = chi-square test
* p<0.05 statistically significant differences between women and men

Table 2: Marital status and gender in elderly from Barra do Garças, MT, Brazil.

| Gender | Marital status | | | | | | Total | |
|--------|----------------|------|-----------|------|---------|------|-------|-------|
| | without bond* | | with bond | | widower | | n | % |
| | n | % | N | % | n | % | n | % |
| Female | 27 | 13.9 | 59 | 30.4 | 108 | 55.7 | 194 | 64.7 |
| Male | 34 | 32.1 | 59 | 55.7 | 13 | 12.3 | 106 | 35.3 |
| Total | 61 | 20.3 | 118 | 39.3 | 121 | 40.3 | 300 | 100.0 |

*p < 0,05, compared with the other two categories.

Discussion

Data from the current study revealed a predominance of a less older population, since the age group of 65-69 years were prevalent (57.2%). These data supports the notion that demographic transition is more recent in Barra do Garças, Mato Grosso State, compared to the cities from the Southeast and Southern Brazil.

In the current work, women (65%) were most prevalent compared to the men. A study in a Public Health Unit from Fortaleza, Ceará (Northeast Brazil), 77.1% of the sample were women [8].

A significant proportion of women without income is very worrying, once this variable (income) is associated with increased health risks.

A work in Khorasan, Iran, clearly demonstrated that illiterate subjects presented higher prevalence of diabetes mellitus, whereas the marital status had not been associated with frequency of that disease [4]. In Stockholm, women with lower education presented 150% increased risk of developing diabetes mellitus [9].

In the current work, the frequency of illiterate men (37.7%) was higher than women (21.6%), which is different from a Brazilian study in Maringá, South Brazil. In that study, 29,2% of men and 24.2% of women were illiterate [10]. Notwithstanding, in the current work the illiteracy (21.4%) was lower compared to a previous study covering one municipality from the North and other from the Northeast Brazil, were illiteracy was 69.3% [11].

A greater frequency of illiterate men was related to increased risk of both type 2 diabetes and cardiovascular diseases, among other chronic non-transmissible diseases [12]. There was an inverse correlation between education level and risk of hypertension, and the population belonging to the highest education levels had 13% to 26% decreased risk of that disease [13]. An Indian study, with 19,969 subjects demonstrated that prevalence of hypertension (33.8%) and smoking (56.6%) were higher among those who had lowest education levels compared to the population of the highest education levels (22.7% and 12.5%, respectively) [14].

A study in Bom Jardim de Goiás which is nearby from Barra do Garças revealed that women had better knowledge on diabetes mellitus than men [15].

Another important variable is the socioeconomic level of the elderly population. In this study, 68% of the elderly belonged to the less fortune socioeconomic classes (D and E). It well known that education level influences the socioeconomic status. Belonging to a low

socioeconomic class increased in 170% and 190% the risk of diabetes mellitus among middle-age women and men, respectively [9].

Regarding marital status, 55.7% of women and 12.3% of men were widower. This result did not substantially differ from observed in a study in the municipality of Encruzilhada, inner Rio Grande do Sul state, South Brazil. In that study, 62% of the elderly were widower, comprising 80% of women, and 20% of men [5].

However, our results are different of data found by Pandey [16], in India. That author found a high proportion of married elderly (75.4%); and the divorced/separated elderly presented worst health status compared to current married or never married. By the other side, the present study was more close to the reality reported by Victor et al. [8], which found 50.5% of elderly with no declared affective bond. In the current study, the majority of elderly lived in loneliness.

In elderly people, loneliness is a risk factor for lesser motivation of feeding. Data of the SOLINUT study in Bordeaux, France, showed 42.6% of elderly with an insufficient caloric intake and 21.3% with protein caloric malnutrition [17].

In this manner, when elderly are exposed to lack of opportunities and social contacts they are more prone to present reduced physical capacity [18], as well as cognitive decay, and increased risk of depressive episodes [19]. Men who die early in general are smokers and living alone [20].

Elderly men not married or living alone had a double risk of cognitive decay according to a Dutch study [21].

It was observed 30.4% of women and 55.7% of men presented affective bonds with partners. The frequency of older people with affective bond was minor than that observed by Feliciano et al. [22], which reported 75.4% and 49.8% of men and women, respectively, had affective bond.

In the current study, 27.8% and 32.1% of elderly women and men, respectively, lived alone. These frequencies were higher compared to the reference values for Brazilian population (men = 5.2%; women = 9.66%) [2], in the same manner, the frequency of loneliness in elderly reported here was higher compared to the 11.7% observed by Victor et al. [8] in Fortaleza, Brazil, and the 9.8% found in two municipalities of the North and Northeast Brazil [11]. The data of the further study corroborate the study of Feliciano et al. [22].

It should be noted that loneliness and absence of affective bonds in older age affects both physical and mental health outcomes [6,23]. In the same manner, being widowed was found to be a risk factor for depression according to a recent study [24].

According to the SABE Project in São Paulo, income, education, and female gender were the factors associated with increased risk of difficulties in performing instrumental activities of daily life [25].

Since elderly with preserved physical and cognitive capacities are considered those with a successful aging, social and public health strategies are important to improve social contacts and affective bonds, and avoid social isolation of the elderly.

Conclusion

In the present study elderly people belonged to the lower socioeconomic classes. Although illiteracy of women was lower than the Brazilian reference value, the frequency of male illiteracy reached 38%. A great proportion of those elderly lived alone which provides an increased risk of psychosocial and health problems.

Acknowledgement

To the “Secretaria de Ação Social de Barra do Garças”, MT, Brazil.

References

1. Bernardini-Zambrini DA (2012) Healthy aging and intergenerational solidarity – Latin America and its moment of opportunity. *Colomb Med* 43: 112-113.
2. Instituto Brasileiro de Geografia e Estatística-IBGE (2016) Projection of Brazil's population by age and sex.1980-2050.
3. Ferrari GSL, Barcelos FA, Nery LD, Santos DF dos, Pereira NS, et al. (2012) Idade ativa: qualidade de vida, saúde e recreação em idosos do Brasil. *Rev Iberoamer Salud y Ciudadania* 1: 7-17.
4. Azimi-Nezhad M, Ghayour-Mobarhan M, Parizadeh MR, Safarian M, Esmaeili H, et al. (2008) Prevalence of type 2 diabetes mellitus in Iran and its relationship with gender, urbanisation, education, marital status and occupation. *Singapore Med J* 49: 571-576.
5. Moraes EP de, Rodrigues RAP, Gerhardt TE (2008) Os idosos mais velhos no meio rural: realidade de vida e saúde de uma população do interior gaúcho. *Texto Context Enferm* 17: 374-383.
6. Dahlberga L, McKeel KJ (2014) Correlates of social and emotional loneliness in older people: evidence from an English community study. *Aging Mental Health* 18: 504–514.
7. Jaremka LM, Andridge RR, Fagundes CP, Alfano CM, Potoski SP, et al. (2014) Pain, depression, and fatigue: loneliness as a longitudinal factor. *Health Psychol* 33: 948-957.
8. Victor JF, Ximenes LB, Almeida PC de, Vasconcelos F de F (2009) Perfil sociodemográfico e clínico de idosos atendidos em unidade básica de saúde da família. *Acta Paul Enferm* 22: 49-54.
9. Agardh EE, Ahlbom A, Andersson T, Efendic S, Grill V, et al. (2007) Socio-economic position at three points in life in association with type 2 diabetes mellitus and impaired glucose tolerance in middle-aged Swedish men and women. *International Journal of Epidemiology* 36: 84-92.
10. Meireles VC, Matsuda LM, Coimbra JAH, Mathias TA de F (2007) Características dos idosos em área de abrangência do programa de saúde da família na região noroeste do Paraná: contribuições para a gestão do cuidado em enfermagem. *Saude Soc* 16: 69-80.
11. Cesar JA, Oliveira-Filho JA, Bess G, Cegiela R, Machado J, et al. (2008) Perfil dos idosos residentes em dois municípios pobres das regiões Norte e Nordeste do Brasil: resultados de estudo transversal de base populacional. *Cad Saude Publica* 24: 1835-1845.
12. Gall SL, Abbott-Chapman J, Patton GC, Dwyer T, Venn A (2010) Intergenerational educational mobility is associated with cardiovascular disease risk behaviours in a cohort of young Australian adults: the childhood determinants of adult health (CDAH) study. *BMC Public Health* 10: 55.
13. Conen D, Glynn RJ, Ridker PM, Buring JE, Albert MA (2009) Socioeconomic status, blood pressure progression, and incident hypertension in a prospective cohort study of female health professionals. *Eur Heart J* 30: 1378-1384.
14. Reddy KS, Prabhakaran D, Jeemon P, Thankappan KR, Joshi P, et al. (2007) Educational status and cardiovascular risk profile in Indians. *PNAS USA* 104: 16263-16268.
15. Lemes Dos Santos PF, Dos Santos PR, Ferrari GS, Fonseca GA, Ferrari CK (2014) Knowledge of diabetes mellitus: does gender make a difference? *Osong Public Health Res Perspect* 5: 199-203.



16. Pandey MK (2009) Association between marital status and health: examining the role of age and gender. Delhi, ASARC working paper 10: 18.
17. Ferry M, Sidobre B, Lambertin A, Barberger-Gateau P (2005) The SOLINUT study: analysis of the interaction between nutrition and loneliness in persons aged over 70 years. *J Nutr Health Aging* 9: 261-268.
18. Hajek A, König HH (2016) Longitudinal predictors of functional impairment in older adults in Europe – Evidence from the survey of health, ageing and retirement in Europe. *PLoS ONE* 11: e0146967.
19. Stoeckel KJ, Litwin H (2016) The impact of social networks on the relationship between functional impairment and depressive symptoms in older adults. *Int Psychogeriatr* 28: 39-47.
20. Tell D, Nilsson PM (2006) Early ageing in middle-aged men is associated with adverse social factors and increased mortality risk: the Malmö Preventive Project. *Scand J Public Health* 34: 346-352.
21. Van Gelder BM, Tijhuis M, Kalmijn S, Giampaoli S, Nissinen A, et al. (2006) Marital status and living situation during a 5-year period are associated with a subsequent 10-year cognitive decline in older men: the FINE study. *J Gerontol B Psychol Sci Soc Sci* 61: P213-P219.
22. Feliciano AB, Moraes SA de, Freitas ICM de (2004) O perfil do idoso de baixa renda no município de São Carlos, São Paulo, Brasil: um estudo epidemiológico. *Cad Saude Publica* 20: 1575-1585.
23. Ouakinin S, Barreira DP (2015) Loneliness and mental health. *Acta Méd Port* 28: 130-132.
24. Kabátová O, Puteková S, Martinková J (2016) Loneliness as a risk factor for depression in the elderly. *Clin Soc Work Health Int* 6: 48-52.
25. Santos JL, Lebrão ML, Duarte YA, Lima FD (2008) Functional performance of the elderly in instrumental activities of daily living: an analysis in the municipality of São Paulo, Brazil. *Cad Saude Publica* 24: 879-886.

Copyright: © 2016 Ferrari CKB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Ferrari CKB, Ferrari GSL, Nery LD, dos Santos DF, Pereira NS, et al. (2016) Affective Bond, Loneliness and Socioeconomic Aspects of an Elderly Population in Midwest, Brazil. *Arch Nurs Pract Care* 2(1): 024-027. DOI: <http://dx.doi.org/10.17352/anpc.000009>