

Gyanesh Kumar Sahu<sup>1</sup>, Harish Sharma<sup>2</sup>,  
Anshita Gupta<sup>1</sup>, Chanchal Deep Kaur<sup>1\*</sup>

<sup>1</sup>Shri Rawatpura Sarkar Institute of Pharmacy,  
Kumhari, Durg, Chhattisgarh, India

<sup>2</sup>Shri Shankaracharya Group of Institutions, Faculty  
of Pharmaceutical Sciences, Bhilai, Chhattisgarh,  
India

**Dates:** Received: 28 July, 2015; Accepted: 28  
August, 2015; Published: 31 August, 2015

**\*Corresponding author:** Dr. Chanchal Deep Kaur,  
Associate Professor, Shri Rawatpura Sarkar Institute  
of Pharmacy, Kumhari, Durg, (C.G.), India, Tel: 91-  
9826660819; E-mail: chanchaldeep@rediffmail.com;  
dr.chanchaldeep@gmail.com

[www.peertechz.com](http://www.peertechz.com)

**Keywords:** Microemulsion; Emulsion; Nano  
emulsion; Method of preparation; Pharmaceutical  
Application

## Review Article

# Advancements in Microemulsion Based Drug Delivery Systems for Better Therapeutic Effects

### Abstract

Recent progress in combinatorial drug has led to the generation of a large number of new compounds. microemulsions are versatile systems of great technological and scientific interest to the researchers because of their potential to incorporate a wide range of drug molecules (hydrophilic and hydrophobic) due to the presence of both lipophilic and hydrophilic domains. A micro emulsion is a transparent, thermodynamically stable mixture of two immiscible liquid stabilized by surfactant (or mixture of surfactant). Microemulsions have many advantages for instance, more drug solubility, thermodynamic stability, manufacturing and permeation is easy over conventional formulations that convert them to important drug delivery systems. The design and development of microemulsions aimed at controlling or improving required bioavailability levels of therapeutic agents. Through this review an attempt has been made to focus on several recent developments occurred in the field of microemulsions based applications and which confirms its role as a suitable cargoes for delivery of drugs. In that note, the relevance of this paper and the truncated basic aspects and application on microemulsions are discussed.

## Introduction

Recent progress in combinatorial chemistry has led to the generation of a large number of new compounds. Today, a large percent of these new chemical entities (NCEs) in addition to many existing drugs often show poor solubilization behavior which lead to poor oral bioavailability with wide intra- and inter- subject variation and present formulators with considerable technical challenges. The selection of an appropriate dosage form is critical because a dosage form with poor drug delivery can make a useful drug worthless. Bioavailability has important clinical implications as both pharmacologic and toxic effects are proportional to both dose and bioavailability [1]. A topical preparation pertain to medicaments applied to the surface of a part of the body and is a term used to describe formulations that have effects only in a specific area of the body and are formulated in such a manner that the systemic absorption of the medicament is minimal. The methods involved in conventional topical drug delivery basically involve either assisting or manipulating the barrier function of the skin (topical antibiotics, antibacterial, emollients, sunscreen agents) or breaching the horny layer at the molecular scale so as to direct drugs to the viable epidermal and dermal tissues without using oral, systemic or other therapies [2,3].

## Basic aspects of microemulsion

Micro emulsion is a colloidal dispersion composed of oil phase, aqueous phase, surfactant and co-surfactant at appropriate ratios, which is a single optically isotropic and thermodynamically stable liquid solution with a droplet diameter usually within the range of 10–100nm [4]. Microemulsions have been widely studied to enhance the bioavailability of the poorly soluble drugs. They offer a cost effective approach in such cases. Microemulsions have very low surface tension and small droplet size which results in high absorption and

permeation. Interest in these versatile carriers is increasing and their applications have been diversified to various administration routes in addition to the conventional oral route. This can be attributed to their unique solubilization properties and thermodynamic stability which has drawn attention for their use as novel vehicles for drug delivery [5-7].

Microemulsions have advantages over both colloidal systems under investigation and conventional emulsions, suspensions and micellar solutions and may provide alternative drug carriers [8]. They are promising delivery systems which allow sustained or controlled drug release for percutaneous, peroral, topical, transdermal, ocular and parenteral administration of medicaments. They offer the advantage of spontaneous formation, ease of manufacturing and scale-up, thermodynamic stability, improved drug solubilization of hydrophobic drugs and bioavailability. Also, microemulsions that have inverse micellar structure may be less comedogenic than either creams or solutions [9,10].

Microemulsions are quaternary systems composed of an oil phase, a water system, surfactants and a cosurfactant [11]. These spontaneously formed systems possess specific physicochemical properties such as transparency, optical isotropic, low viscosity and thermodynamic stability. The observed transparency of these systems is due to the fact that the maximum size of the droplets of the dispersed phase is not larger than one-fourth of the wavelength of visible light approximately 150 nm. Droplet diameter in stable microemulsions is usually within the range of 10-100 nm (100-1000 °A), which means that the term 'micro emulsion' is misleading and these systems are actually Nano-sized emulsions. Many studies have shown that micro emulsion formulations possessed improved transdermal and dermal delivery properties, mostly *in vitro* [12-21] and several *in vivo* [22-24].

## Types of microemulsion

Three types of microemulsions (Figure 1) are most likely to be formed depending on the composition:

1. Oil in water (O/W) microemulsions wherein oil droplets are dispersed in the continuous aqueous phase.
2. Water in oil (W/O) microemulsions wherein water droplets are dispersed in the continuous oil phase.
3. Bi-continuous microemulsions wherein micro domains of oil and water are inter dispersed within the system.

In all three types of microemulsions, the interface is stabilized by an appropriate combination of surfactants and co-surfactants [25].

## Comparison between emulsion, nanoemulsion and microemulsion

Emulsions and microemulsions are both stable dispersions of oil-in-water or water-in-oil. In emulsion systems, the structures are large enough to scatter light and as such they appear as cloudy colloidal solutions in comparison. The gross physical differences between micro emulsion and emulsion systems can be determined by visual examination- microemulsions show no tendency to phase separate and are usually optically transparent, whereas emulsions are opalescent or turbid and the phases inevitably separate (Table 1).

## Formulation considerations

The challenges in formulating microemulsions are (Figure 2):

1. Determining systems that are non-toxic, non-irritating, non-comedogenic and non-sensitizing.

2. Formulating cosmetically elegant microemulsions. The micro emulsion formulation must have low allergic potential, good physiological compatibility and high biocompatibility. The components involved in the general formulation of microemulsions include (a) an oil phase (b) an aqueous phase containing hydrophilic active ingredients [preservatives and buffers may be included] (c) a primary surfactant [anionic, non-ionic or amphoteric] (d) secondary surfactant or cosurfactants.

## Construction of phase diagram

Pseudo-ternary phase diagram is constructed to obtain the appropriate components and their concentration ranges that can result in large existence area of micro emulsion. Once the appropriate micro emulsion components are selected, ternary pseudo phase diagram is constructed to define the extent and nature of the micro emulsion regions. To produce such diagrams, a large number of samples of different composition are prepared [31-39]. To study the phase behavior of simple micro emulsion systems comprising of surfactant, oil and water at fixed pressure and temperature ternary phase diagrams are used. Each corner of the ternary phase diagram represents 100% concentrations of a particular component. When four or more components are used pseudo-ternary phase diagrams are used to depict these systems in which each corner represents binary mixtures of two components such as surfactant/co-surfactant, surfactant/water, oil/drug, and water/drug mixtures [40].

## Components of microemulsion formulations

**Water phase:** Depending upon the amount of water present in the system, water may form water pool or work as a dispersion medium in micro-emulsion systems [41].





Figure 1: Different types of microemulsions formed depending on the composition.

**Table 1:** Comparison between emulsion and microemulsion [26-30].

Sr. No.	Specification	Emulsion	Nanoemulsion	Microemulsion
1.	Appearance	Most emulsions are opaque (white) because bulk of their droplets is greater than wavelength of light and most oils have higher refractive indices than water.	Nanoemulsions are part of a broad class of multiphase colloidal dispersions. It is dispersions of nanoscale droplets formed by shear-induced rupturing.	Microemulsions are transparent or translucent as their droplet diameter are less than $\frac{1}{4}$ of the wave length of light, they scatter little light.
2.	Optical Isotropy	Anisotropic	Isotropic	Isotropic
3.	Microstructure	Static	Dynamic	Dynamic
4.	Droplet Size	>500nm	1-100nm	20-200nm
5.	Droplet Shape	Emulsions consist of roughly spherical droplets of one phase dispersed into the other.	Nanoemulsion consist of spherical in shape due to that, small-sized droplet with its high surface area allowing effective transport of the active to the bioactive effects.	They constantly evolve between various structures ranging from droplet like swollen micelles to bicontinuous structure.
6.	Molecular Packing	Inefficient	Efficient	Efficient
7.	Stability	Thermodynamically unstable (Kinetically stable), will eventually phase separate.	Thermodynamically stable, long shelf-life	Thermodynamically stable, long shelf-life
8.	Phase	Biphasic	Monophase	Monophase
9.	Viscosity	High	Low	Low
10.	Interfacial tension	High	Low	Ultra low

**Figure 2:** Schematic representation of formulation of Microemulsion.

**Oil phase:** The oil phase must be chosen appropriately, since it governs the selection of the other ingredients for the micro emulsion and there are two main factors that need be considered before selecting the appropriate oil phase. Firstly, the solubilising potential of the oil for the selected substance must be seen. And secondly, the chosen must be such that the micro emulsion forming region is enhanced. Oils with shorter hydrocarbon chains are easier to micro-emulsify as compared to oils with long hydrocarbon chains. An oils ability to solubilise lipophilic groups is directly proportional to the chain length of the oil. Thus, the selected oil should be such that it is capable of solubilising the API, and facilitating the formation of microemulsions with desired characteristics [42,43].

### Surfactants in microemulsions

Surfactants are molecules that typically contain a polar head group and an apolar tail [44]. They are surface-active and microstructure-forming molecules with a strong chemical dipole [45]. They can be ionic (cationic or anionic), nonionic, or zwitterionic. Surfactant molecules self-associate due to various inter- and intra-molecular forces as well as entropy considerations. The surfactant molecules can arrange themselves in a variety of shapes. They can form spherical micelles, rod-shaped micelles, a hexagonal phase (consisting of rod-shaped micelles), lamellar (sheet) phases, reverse micelles, or hexagonal reverse micelles [46].

### Cosurfactants

In most cases, single-chain surfactants alone are unable to reduce the o/w interfacial tension sufficiently to enable a micro emulsion to form [47-50]. The presence of cosurfactants allows the interfacial film sufficient flexibility to take up different curvatures required to form micro emulsion over a wide range of composition. If a single surfactant film is desired, the lipophilic chains of the surfactant should be sufficiently short, or contain fluidizing groups (e.g. unsaturated bonds). Short to medium chain length alcohols (C3-C8) are commonly added as cosurfactants which further reduce the interfacial tension and increase the fluidity of the interface [51-53] (Table 2).

### Applications of microemulsion

During the last two decades, microemulsions have been extensively researched because of their tremendous potential in many

applications. The role of microemulsions in various field are (Figure 3).

### Microemulsion in pharmaceutical

**Parenteral application:** Parenteral administration (especially via the intravenous route) of drugs with limited solubility is a major problem in the pharmaceutical industry because of the extremely low amount of drug actually delivered to a targeted site. Micro emulsion formulations have distinct advantages over macro emulsion systems when delivered parenterally because of the fine particle, micro emulsion is cleared more slowly than the coarse particle emulsion and, therefore, have a longer residence time in the body. Both O/W and W/O micro emulsion can be used for parenteral delivery [55]. Rhee et.al formulated itraconazole containing parentral micro emulsion, using an o/w micro emulsion system. he average droplet size of the microemulsions was < 150 nm, and the hemolysis test showed this formulation to be nontoxic to red blood cells. The pharmacokinetic profiles of the ITZ-micro emulsion for itraconazole and its major metabolite, hydroxyitraconazole, were compared with those of a PEG 400 solution and cyclodextrin formulations in rats. Overall, these results highlight the potential of an ITZ-micro emulsion formulation for the parenteral route [56].

### Oral administration


Oral administration of micro emulsion formulations offer the several benefits over conventional oral formulation for oral administration including increased absorption, improved clinical potency, and decreased drug toxicity. Therefore, micro emulsion has been reported to be ideal delivery of drugs such as steroids, hormones, diuretic and antibiotics [56].

The development of the effective oral delivery systems has always been the main goal because drug efficacy can be severely limited by instability or poor solubility in the gastrointestinal fluid. Biopharmaceutical Classification System (BCS) is a useful guidance by US FDA shown in Table 3 and it takes into account contributions of three major factors, dissolution, solubility, and intestinal permeability, which affect oral drug absorption.

microemulsions have the potential to enhance the solubilization of the poorly soluble drugs and overcome the dissolution related bioavailability problems. This is particularly important for the BCS

**Table 2:** Some commonly used components for microemulsions [54].

Components	Examples	
Oil	MCTs	Glyceryl tricaprlylate/caprante: Captex 355, Miglyol 810, Neober M-5 etc
	LCTs	Corn oil, soyabean oil, safflower oil, olive oil etc
	Mono/diglyceride	Glyceryl caprylate/caprante (Capmul MCM), Glycerol monooleate (Capmul GMO) etc
	Fatty acids	Oleic acid
	Propylene glycol ester	Capmul PG-8, Propylene glycol monolaurate (Lauroglycol)
Surfactant	HLB > 10	Tween-20, Tween-80, Polyoxyl 35 castor oil (Cremophore EL), PEG-8 caprylic/ capric glycerides (Labrasol), Polyoxyl 40 hydrogenated castor oil (Cremophore RH 40) etc.
	HLB < 10	Phosphatidylcholine, Unsaturated polyglycolized glycerides (Labrafil M 2125), Span-40, Span-80 etc
Co-surfactant	Propylene glycol, Polyethylene glycol, Ethanol, Isopropyl alcohol, Isopropyl myristate, ethanol, propanol, isopropanol, etc.	



**Table 3:** According to the BCS, classification of drug substances.

Sr. No.	Class	Permeability	Solubility
1.	I	High	High
2.	II	High	Low
3.	III	Low	High
4.	IV	Low	Low

class II or class IV drugs. The successful formulation of such drugs is highly dependent on the performance of the formulated product. microemulsions act as super solvent of these drugs and can be optimized to ensure consistent bioavailability. In addition, they can be used for the delivery of hydrophilic drugs including macromolecules such as proteins and peptides. This is due to the existence of polar, nonpolar and interfacial domains which allow encapsulation of drugs with varying solubility [57,58].

### Topical administration

Topical administration of drugs can have advantages over other methods for several reasons, one of which is the avoidance of hepatic first pass metabolism of the drug and related toxicity effects. Another is the direct delivery and targetability of the drug to affected area of the skin or eyes [58].

Due to the small micelles size and large amount of inner phase in microemulsions, the density of droplets and their surface area are

assumed to be high. Therefore, micelles settle down to close contact with the skin providing high concentration gradient and improved drug permeation. Moreover, low surface tension ensures good contact to the skin. Also, the dispersed phase can act as a reservoir making it possible to maintain an almost constant concentration gradient over the skin for a long time [59].

**Nasal delivery:** Microemulsions are now being studied as a delivery system to enhance uptake across nasal mucosa. Addition of a mucoadhesive polymer helps in prolonging the residence time on the mucosa [60,61]. Sheetal et al. in 2013, Prepared and evaluated transnasal micro emulsion of carbamazepine. Oleic acid was selected as oil while Tween 80 and propylene glycol were selected as surfactant and cosurfactant respectively based on solubility results. Optimized ratio of Tween 80: propylene glycol was selected after developing pseudoternary phase diagrams for different ratio and microemulsions were prepared. The prepared microemulsions were evaluated for globule size, viscosity, pH, conductivity and % transmittance. Ex-vivo diffusion study for optimized micro emulsion was performed through sheep nasal mucosa wherein diffusion flux and permeability coefficients were determined. Further pharmacodynamics performance was evaluated in rats by electrically induced seizures. It was found that optimized micro emulsion was stable and transparent with average globule size of 190 nm and diffusion flux of  $75.77 \mu\text{g cm}^{-2} \text{min}^{-1}$  and showed no toxicity during histopathological evaluation on sheep nasal mucosa.

## Ophthalmic delivery

In conventional ophthalmic dosage forms, water soluble drugs are delivered in aqueous solution while water insoluble drugs are formulated as suspensions or ointments. Low corneal bioavailability and lack of efficiency in the posterior segment of ocular tissue are some of the serious drawbacks of these systems. microemulsions have emerged as a promising dosage form for ocular useful in achieving improved patient compliance and favorable for ophthalmological use [62]. Okur et al. 2014, studied the effect of Novel ofloxacin-loaded micro emulsion formulations for ocular delivery and found that OFX microemulsions could be offered as a promising strategy for ocular drug delivery.

## Tumor targeting

The utility of microemulsions as vehicles for the delivery of chemotherapeutic or diagnostic agents to neoplastic cells while avoiding normal cells. A method for treating neoplasms, wherein neoplasms cells have an increased number of LDL (low density, lipoprotein) receptors compared to normal cells. The micro emulsion comprised of a nucleus of cholesterol esters and not more than 20% triglycerides surrounded by a core of phospholipids and free cholesterol and contained a chemotherapeutic drug. The microemulsions could then be incorporated into cells via receptors for LDL and delivered the incorporated molecules. Thus, higher concentration of anticancer drugs could be achieved in the neoplastic cells that have an increased expression of the receptors. In this way toxic effects of these drugs on the normal tissues and organs could be avoided [69,70].

## Microemulsions in cosmetics

In many cosmetic applications such as skin care products,

emulsions are widely used with water as the continuous phase. It is believed that micro emulsion formulation will result in a faster uptake into the skin. Cost, safety (as many surfactants are irritating to the skin when used in high concentrations), appropriate selection of ingredients (i.e. surfactants, cosurfactants, oils) are key factors in the formulation of microemulsions [77,78].

## Solubilization of drug in microemulsion

Microemulsion possesses interesting physicochemical properties, i.e. transparency, low viscosity, thermodynamic stability, high solubilization power. Because of these specific properties of micro emulsion can be useful as a drug delivery system. The different categories of drugs solubilized in micro emulsion systems for their better therapeutic efficacy [79] (Table 4).

## Conclusion

Microemulsions are an attractive technology platform for the pharmaceutical formulators as it has excellent transparency and relatively simple formulation process. The role of micro emulsion systems is of paramount importance in providing novel solutions to overcome the problems of poor aqueous solubility of highly lipophilic drug compounds and provide high, more consistent and reproducible bioavailability. Microemulsions have proved their efficiency in oral administration of cyclosporine (Neoral). But the main concern is that the applicability and effectiveness of micro emulsion based drug delivery systems should not be limited to oral routes only. They can be easily used for developing novel cargoes for phytoactives and extracts also. The current scenario potentiate the targeting aspects of microemulsions due to their tremendous capability of carrying lipophilic drugs. Several targeting cargoes have been developed in recent years with the concept of microemulsions at its base. It can be hoped that in near future microemulsions will

**Table 4:** Research envisages of microemulsion in different route of administration.

Sr. No.	Delivery System	Drug	Category	Application	Ref.
1.	Nasal Delivery	Diazepam	Anticonvulsant or antiepileptic drug	Nasal route for administration of diazepam is useful approach for the rapid onset of action during the emergency treatment of status epilepticus.	[61]
2.	Ophthalmic Delivery	Dexamethasone	Antiallergic	It showed better tolerability and higher bioavailability. The formulation showed greater penetration in the eye which allowed the possibility of decreasing the number of applications per day.	[63]
3.	Parenteral Application	vitamins E, A, D, and K	Supplements	It suitable for fat soluble vitamins and hydrophobic drugs	[64]
4.	Oral administration	Paclitaxel	Anticancer	Microemulsion permitted its rapid and efficient absorption resulting in improved oral bioavailability.	[65]
5.	Topical administration	miconazole, ketoconazole, itraconazole	Antimycotics	microemulsions impart to them increased drug loading, enhanced penetration through the biological membranes, and increased bioavailability.	[66,67]
6.	Tumor targeting	Aclacinomycin	Antitumor agent	Folate-linked microemulsion is feasible for tumour targeted ACM delivery. The study showed that folate modification with a sufficiently long PEG chain on emulsions is an effective way of targeting tumour cells.	[68]
7.	Brain targeting	Clonazepam	Anticonvulsant or antiepileptic drug	Mucoadhesive microemulsion compared to i.v. was found to be 2-fold higher indicating larger extent of distribution of the drug in the brain.	[76]
8.	Cosmetic	-	Moisturizing, soothing agents, sunscreens, antiperspirants, body cleansing agents, hair conditioners.	They are now being widely investigated for preparing personal care products with superior features such as having improved product efficiency, stability, appearance and minimal irritation.	[79]

become a versatile system for drug delivery not only for conventional but also for phytoactive and that the potential of microemulsions as novel compartmentalized liquids will be even more significant in upcoming era.

## References

- Aungst BJ (1993) Novel formulation strategies for improving oral bioavailability of drugs with poor membrane permeation or presystemic metabolism. *J Pharm Sci* 82: 979-987.
- Gundogdu E, Baspinar Y, Koksai C (2013) A Microemulsion for the Oral Drug Delivery of Pitavastatin. *Pharmaceut Anal Acta* 4: 1-5.
- Bidyut KP, Satya PM (2001) Uses and applications of microemulsions. *Current Science* 80: 990-1001.
- Changez M, Varshney M (2000) Aerosol-O microemulsions as transdermal carriers of tetracaine hydrochloride. *Drug Dev. Ind. Pharm* 26: 507-512.
- Kovarik JM, Mueller EA, van Bree JB, Tetzloff W, Kutz K (1994) Reduced inter and intra individual variability in cyclosporine pharmacokinetics from a microemulsion formulation. *J Pharm Sci* 83: 444-446.
- Noble S, Markham A (1995) Cyclosporin. A review of the pharmacokinetic properties, clinical efficacy and tolerability of a microemulsion-based formulation (Neoral). *Drugs* 50: 924-941.
- Erkko P, Granlund H, Nuutinen M (1997) Comparison of cyclosporine A pharmacokinetics of a new microemulsion formulation and standard oral preparation in patients with psoriasis. *Br J Dermatol* 136: 82-88.
- Jain NK (2010) Progress in controlled and novel drug delivery systems. 1st ed, New Delhi: CBS publishers and distributors 324.
- Moghimpour E, Salimi A, Eftekhari S (2013) Design and Characterization of Microemulsion Systems for Naproxen. *Adv Pharm Bull* 3: 63-71.
- Khalil E, Shorouq T, Al-Sotari (2012) Formulation and Characterization of IPM/Water/Nonionic-Ionic Surfactant Microemulsions. *J Chem Chem Eng* 6: 187-198.
- Ceglie A, Das KP, Lindman B (1987) Effect of oil on the microscopic structure in four component cosurfactants microemulsions. *Colloids Surf* 115: 115-120.
- Osborne DW, O'Neill AJ (1991) Microemulsions topical drug delivery vehicles: in-vitro transdermal studies model hydrophilic drug. *J Pharm Pharmacol* 43: 450-454.
- Trotta M, Pattarino F, Gasco MR (1996) Influence of counter ions on the skin permeation of methotrexate from water-oil microemulsions. *Pharm. Acta Helv* 71: 135-140.
- Delgado-Charro MB, Iglesias-vilasG, Blanco-Mendez J et al. (1997) Delivery of hydrophilic solute through then skin from novel microemulsion system. *Eur J Pharm Biopharm* 43: 37-42.
- Parra JL, Coderch L, Yuste I, A de la Maza (1997) Incorporation of non-steroidal anti-inflammatory drugs into specific monophasic formulation. *Colloids Surf A: Physicochem Eng. Aspect* 123-124: 115-123.
- Dreher F, Walde P, Walther P, Wehrli E (1997) Interaction of a Lecithin Microemulsion Gel with Human Stratum Corneum and its Effect on Transdermal Transport. *J Control Release* 45, 131-40.
- Schmalfuss U, Neubert R, Wohlarb W (1997) Modification of drug penetration into human skin using microemulsion. *J Control Release* 46: 279-285.
- Kreilgaard M, Pedersen E J, Jaroszewski JW (2000) NMR characterization and transdermal drug delivery potential of microemulsion systems. *J Control Release* 69: 421-33.
- Alvarez-Figueroa M J, Blanco-Mendez J (2001) Transdermal delivery of methotrexate: iontophoretic delivery from hydrogels and passive delivery from microemulsions. *Int J Pharm* 215: 57-65.
- Rhee YS, Choi JG, Park ES, Chi SC (2001) Transdermal delivery of ketoprofen using microemulsions. *Int J Pharm* 228: 161-170.
- Lee PJ, Langer R, Shastri VP (2003) Novel microemulsion enhancer formulation for simultaneous transdermal delivery of hydrophilic and hydrophobic drugs. *Pharm Res* 20: 264-269.
- Kemken J, Ziegler A, Mueller BW (1992) Influence of supersaturation on the pharmacodynamic effect of Bupranolol after dermal administration using microemulsions as vehicle. *Pharm Res* 9: 554-58.
- Kreilgaard M (2001) Dermal pharmacokinetics of microemulsion formulations determined by in-vitro microdialysis. *Pharm. Res* 18: 367-373.
- Singh MK, Chandel V, Gupta V (2010) Formulation development and characterization of microemulsion for topical delivery of Glipizide. *Scholars Res Lib* 3: 33-42.
- Schulman JH, Stoekenius D, Prince LM (1959) Mechanism of formation and structure of microemulsions by electron microscopy. *J Phy Che* 63: 1677-1680.
- Kayes FB (1999) Disperse systems. In Aulton ME (Ed.) *Pharmaceutics: The science of dosage form design, international student edition*. Churchill Livingstone 110.
- Rieger MM (1987) Emulsions. In Lachman L, Lieberman HA, Kanig JL *Theory and practice of industrial pharmacy*. 3rd ed. Varghese Publishing House, Bombay 507-519.
- Eccleston M (2002) Emulsion and microemulsions. In Swarbrick J, Boylan JC (ed.) *Encyclopedia of pharmaceutical technology*. 2nd edition. Marcel Dekker, Inc., New York 2: 1080-1085.
- Betageri G, Prabhu S (2002) Semisolid preparations. In Swarbrick J, Boylan JC (ed.) *Encyclopedia of pharmaceutical technology*. 2nd edition. Marcel Dekker, Inc., New York 3: 2441- 2442.
- Ghosh PK, Murthy RSR (2006) Microemulsions: A potential drug delivery system. *C Drug Del* 3: 167-80.
- Cui F, Qian F, Yin C (2006) Preparation and characterization of mucoadhesive polymer-coated nanoparticles. *Int J Pharmaceutics* 316, 1-2: 154-161.
- Sharma G, Mishra AK, Mishra P, Misra A (2009) Intranasal Cabergoline: Pharmacokinetic and Pharmacodynamic Studies. *AAPS Pharm Sci Tech* 10: 1321-1330.
- Kaler EW, Prager S (1982) A model of dynamic scattering by microemulsions. *J Colloid Interface Sci* 86: 359-369.
- Lawrence JM, Rees GD (2000) Microemulsion based media as novel drug delivery systems. *Adv Drug Del Rev* 45: 89-121.
- Lianli L, Nandi I, Kim KH (2002) Development of an ethyl laurate-based microemulsion for rapid-onset intranasal delivery of diazepam. *Int J Pharmaceutics* 237: 77-85.
- Roland I1, Piel G, Delattre L, Evrard B (2003) Systemic characterization of oil-in-water emulsions for formulation design. *Int J Pharm* 263: 85-94.
- Faiyaz Shakeel, Sanjula Baboota, Javed Ali, Mohammed Aqil, Sheikh Shafiq et al. (2007) Nano emulsions as vehicles for transdermal delivery of Aceclofenac. *AAPS Pharm Sci Tec* 8: 191-199.
- Ugwoke MI, Verbeke N, Kinget R (2001) The biopharmaceutical aspects of nasal mucoadhesive drug delivery. *J Pharm Pharmacol* 53: 13-21.
- Zhang Q, Jiang X, Jiang W, Lu W, Su L, et al. (2004) Preparation of nimodipineloading microemulsion for intranasal delivery and evaluation of the targeting efficiency to brain. *Int J Pharm* 275: 85-96.
- Shaji JR (2004) Microemulsions as drug delivery systems. *Pharma Times* 36: 17-24.
- Chaparaba DP (1991) Studies on the properties of normal and reversed micellar systems. PhD thesis submitted in Department of Chemistry, Aligarh Muslim University, Aligarh, India 64.
- Rana S, Verma G, Hassan PA (2013) Microstructural investigation of lipid

- solubilized microemulsions using laser light scattering. *Adv. Mat. Lett* 4: 476-481.
43. Warisnoicharoen W, Lansley AB, Lawrence MJ (2000) Nonionic oil in water microemulsion: The effect of oil type on phase behavior. *Int J Pharm* 198: 7-27.
44. Evans DF, Wennerstrom H (1999) *The Colloidal Domain. Where Physics, Chemistry, and Biology Meet*, 2nd ed, Wiley-VCH, New York 632.
45. Holmberg K, Shah DO, Schwuger MJ (2002) *Handbook of applied surface and colloid chemistry*. Chichester, New York 591.
46. Lawrence MJ, Rees GD (2002) Microemulsion-based media as novel drug delivery systems. *Advanced Drug Delivery Reviews* 45: 89-121.
47. Bhargava HN, Narurkar A, Lieb LM (1987) Using microemulsions for drug delivery. *Pharm Tech* 11: 46-52.
48. Kreuter J (1994) Microemulsions, In: *Colloidal drug delivery systems*. New York: Marcel Dekker 31-71.
49. Lawrence MJ (1984) Surfactant systems: microemulsions and vesicles as vehicles for drug delivery. *Eur J Drug Metab Pharmacokinet* 3: 257-269.
50. Tenjarla S (1999) Microemulsions: an overview and pharmaceutical applications. *Crit Rev Therapeut Drug Carrier Syst* 16: 461-521.
51. Ghosh PK, Murthy RSR (2006) Microemulsions: A potential drug delivery system. *Curr Drug Deliv* 3: 167-180.
52. Aboofazeli R, Lawrence CB, Wicks SR et al. (1994) Investigations into the formation and characterization of phospholipid microemulsions III. Pseudo-ternary phase diagrams of systems containing water-lecithin-isopropyl myristate and either an alkanolic acid, amine, alkanediol, poly ethylene glycol alkyl ether or alcohol as co-surfactant. *Int J Pharm* 111: 63-72.
53. Stilbs P, Lindman B, Rapacki K (1983) Effect of alcohol cosurfactant length on microemulsion structure. *J Colloid Interface Sci* 95:583-585.
54. Cannon JB (2010) Lipid-based Formulatin Approaches for Poorly Soluble drugs
55. Ho H, Hsiao CC, Sheu MT (1996) Preparation of microemulsions using polyglyceryl fatty acid esters as surfactant for the delivery of protein drugs. *J Pharm Sci* 85: 138-143.
56. Rhee YS, Park CW, Nam TY, Shin YS, Chi SC, et al. (2007) Formulation of parenteral microemulsion containing itraconazole. *Arch Pharm Res* 30: 114-123.
57. Sharma H, Kumar Sahu G, Dapurkar V (2012) An overview of new drug delivery system: microemulsion. *Int. Res J Pharm App Sci* 2: 1-8.
58. Sariciaux MJ, Alan L, Sado PA (1995) Using microemulsion for drug delivery of therapeutic peptides. *Int J Pharm* 1995, 12:127-36.
59. Swenson EC, Curatolo WJ (1992) Intestinal permeability enhancement for proteins, peptides and other polar drugs: mechanism and potential toxicity. *Adv Drug Del Rev* 8: 39-42.
60. Ho H, Huang MC, Chen LC (1998) The percutaneous delivery of prostaglandin E1 and its alkyl esters by microemulsions. *Chin Pharm J* 50: 257-266.
61. Sheetal Porecha Acharyaa, K. Pundarikakshudu, Aashish Panchal, Anita Lalwani, (2013) Preparation and evaluation of transnasal microemulsion of carbamazepine; *Asian Journal of Pharmaceutical Sciences* 8: 64-70.
62. Lianly IL, Nandi I, Kim KH (2002) Development of an ethyl laurate based microemulsion for rapid onset of intranasal delivery of diazepam. *Int J Pharm* 237: 77-85.
63. Üstündag-Okur N, Gökçe EH, Eerilmez S, Özer Ö, Ertan G (2014) Novel ofloxacin-loaded microemulsion formulations for ocular delivery. *J Ocul Pharmacol Ther* 30: 319-332..
64. Fialho SL, Da Silva-Cunha A (2004) New vehicle based on a microemulsion for topical ocular administration of dexamethasone. *Clin Expt Opthal* 32: 626-632.
65. Vigne JL, Kane JP (1991) US5023271.
66. Gao P, Morozowich W (2006) US20067115565.
67. Tenjarla SN (1999) Microemulsions: An overview and pharmaceutical applications. *Crit Rev Ther Drug Carrier Syst* 16: 461-521.
68. Lieberman HA, Rieger MM, Banker GS (1996) *Pharmaceutical Dosage Forms, Disperse Systems*. 2nd ed. Vol 1. New York: Marcel Dekker Inc 211-281: 315-370.
69. Shiokawa T, Hattori Y, Kawano K, Ohguchi Y, Kawakami H et al. (2005) Effect of polyethylene glycol linker chain length of folate-linked microemulsions loading aclacinomycin A on targeting ability and antitumour effect in vitro and in vivo. *Clin Cancer Res* 11: 2018-2025.
70. Maranhao RC (1996) US5578583.
71. Shiokawa T, Hattori Y, Kawano K, Ohguchi Y, Kawakami H et al. (2005) Effect of polyethylene glycol linker chain length of folate-linked microemulsions loading aclacinomycin A on targeting ability and antitumour effect in vitro and in vivo. *Clin Cancer Res* 11: 2018-2025.
72. Wermeling DP, Miller JL, Archer SM, Manaligod JM, Rudy AC (2001) Bioavailability and pharmacokinetics of lorazepam after intranasal, intravenous and intramuscular administration. *J Clin Pharmacol* 41: 1225-1231.
73. Dorman DC, Brenneman KA, McElveen AM, Lynch SE, Roberts KC, et al. (2002) Olfactory transport: A direct route of delivery of inhaled manganese phosphate to the rat brain. *J Toxicol Environ Health* 65: 1493-1511.
74. Draghia R, Caillaud C, Manicom R, Pavirani A, Kahn A, et al. (1995) Gene delivery into the central nervous system by nasal instillation in rats. *Gene Ther* 2: 418-423.
75. Illum L (2000) Transport of drugs from the nasal cavity to central nervous system. *Eur J Pharm Sci* 11: 1-18.
76. Talegaonkar S, Mishra P (2004) Intranasal delivery: An approach to bypass the blood brain barrier. *Ind J Pharmacol* 36: 140-147.
77. Vyas SP (2009) *Theory and practice in novel drug delivery system*. CBS Publishers, New Delhi 115-116.
78. Shinoda K, Shibata Y, Lindman B (1993) Interfacial tensions for lecithin microemulsions including the effect of surfactant and polymer addition. *Langmuir* 9: 1254-1257.
79. Azeem A, Rizwan M, Ahmad FJ, Khan ZI, Khar RK, et al. (2008) Emerging role of microemulsions in cosmetics. *Recent Pat Drug Deliv Formul* 2: 275-289.
80. Winsor PA (1954) *Solvent properties of amphiphilic compounds*. Butterworth, London 207.

**Copyright:** ©2015 Sahu GK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Sahu GK, Sharma H, Gupta A, Kaur CD (2015) Advancements in Microemulsion Based Drug Delivery Systems for Better Therapeutic Effects. *Int J Pharm Sci Dev Res* 1(1): 008-015.