



## Short Communication

# Pre-symptomatic and asymptomatic COVID-19 cases in Nigeria amidst prevailing socio-cultural beliefs and practices: Implication for COVID-19 transmission and way forward

Elvis Efe Iser<sup>1\*</sup>, Adewale Moses Adejgbagbe<sup>1</sup>, Victoria Oladoyin<sup>2</sup>, Samuel Abiona<sup>3</sup> and Eniye Nosa Omorogbe<sup>4</sup>

<sup>1</sup>University of Ibadan, Ibadan, Nigeria

<sup>2</sup>University of Medical Sciences, Ondo, Nigeria

<sup>3</sup>Ondo State Primary Health Care Development Agency, Akure, Nigeria

<sup>4</sup>Medical Laboratory Science Council of Nigeria, Abuja, Nigeria

**Received:** 08 August, 2020

**Accepted:** 24 August, 2020

**Published:** 25 August, 2020

**\*Corresponding author:** Elvis Efe Iser, Department of Epidemiology and Medical Statistics, Faculty of Public Health, College of Medicine, University of Ibadan, Ibadan, Nigeria, Tel: +2348030480305; E-mail: [elvisisere@gmail.com](mailto:elvisisere@gmail.com), [iserielvisevdestiny@yahoo.co.uk](mailto:iserielvisevdestiny@yahoo.co.uk)

**Keywords:** Pre-symptomatic; Asymptomatic; COVID-19; Socio-culture beliefs; Health behaviour

<https://www.peertechz.com>



## Abstract

Nigeria is the most populous country in Africa with diverse range of cultural and religious beliefs and practices vis-à-vis health seeking behaviour. The country is currently one of the nations in the African region that is most affected by the COVID-19 pandemic. With the active outbreak of COVID-19 in the country, this paper discussed how key socio-cultural beliefs and health behaviour coupled with pre-symptomatic and asymptomatic clinical presentation of COVID-19 cases could foster the spread of the disease and recommendations for control.

## Introduction

COVID-19, a viral disease caused by the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), has spread rapidly over multiple countries in all the continents of the world since the first localized outbreak in China in December, 2019 with a resultant considerable impact on global morbidity and mortality [1]. Currently, Nigeria ranked second among the countries within the African region with ongoing transmission of COVID-19 disease [2]. As at July 31<sup>st</sup>, 2020, Nigeria has recorded a total of 43, 151 confirmed cases and 879 deaths in all 36 states and Federal Capital Territory since the first case was confirmed in the country on February 27, 2020 [3].

The COVID-19 outbreak which has been declared as a public health emergency of international concern by the World Health Organization (WHO) is exponentially sweeping across the world and triggering chaos, fear, anxiety and stress in countries [4-7]. The pandemic has disrupted daily routine lifestyles and business, stock market and global educational system. Similarly, the consequences of the COVID-19 pandemic in Nigeria have resulted in economic lockdown of major cities, total shutdown of the educational system across the country and economic and physical hardship among the populace [4,8].

The incubation period for COVID-19, which is the time between exposure to the virus and onset of symptom(s), is



on average 5–6 days but can be up to 14 days [9]. During this period, also known as the pre-symptomatic period, COVID-19 transmission has been documented through contact tracing efforts and enhanced investigation of clusters of confirmed cases [10–15]. Data suggesting that some people can test positive for COVID-19 from 1–3 days before they develop symptoms also supports pre-symptomatic transmission of COVID-19 [14,16]. Given this background, it is important to note that pre-symptomatic cases, that are persons who may not have symptoms at the early stage of the illness but who will later develop symptoms, can transmit the virus before significant symptoms develop via infectious droplets [2,14].

Similarly, preliminary findings from the field are suggestive that asymptomatic cases, defined as persons with COVID-19 who do not have symptoms and never develops symptoms, have the capacity to infect others within close contact [17,18].

The pre-symptomatic and asymptomatic COVID-19 cases are active silent carriers of the virus who could infect other persons they come in close contact with unknowingly resulting in super-spreading of the disease where a single patient could infect a disproportionate number of contacts and likely contribute to the speed and degree of the outbreak [14,19].

Epidemiological review of the COVID-19 outbreak in Nigeria has shown laboratory-confirmed cases that are either pre-symptomatic or asymptomatic [20].

### **Socio-cultural beliefs/practices and health seeking behaviour in Nigeria: key concerns**

Nigeria, a country made up of people from different ethnic groups, has more than 250 ethnic groups with different cultural and religious beliefs. These beliefs are governed by societal norms based on age, gender, religion, social class that could influence the perception, attitudes and behaviour of individuals towards disease existence, risk perception, etiology, severity and treatment [21,22].

As in many other developing countries, some ethnic groups in Nigeria have continued to link the etiology of diseases with spiritual or supernatural and traditional concepts despite medical advances which have led to improved understanding of the aetiology of diseases [22,23]. A typical example is the cultural belief among some ethnic groups that illnesses are as a result of the direct actions of supernatural entities, such as gods, evil or ancestral spirits and thus demands spiritual and traditional solution [23].

Furthermore, as a result of the cultural and religious beliefs in the Nigerian context, a lot of people are optimistic about their health and hence perceive themselves as having a low risk of falling ill [18]. The health belief model however predicts that individuals with low self-perceived risk of becoming sick are more likely to engage in unhealthy or risky behaviours [24].

Another socio-cultural practice in the country is the preference for non-orthodox care, such as visiting traditional healers and taking herbal concoctions, when ill because of the belief that traditional healers are easily accessible compared

to visiting a medical personnel in a hospital; and also because of the belief that herbal remedies are readily available, less expensive, more efficient and with less adverse effect compared with orthodox drugs. Other socio-cultural practices in Nigeria include engagement in mass religious gatherings due to the belief that healing and good health depends on the intervention of the supreme God as well as engagement in traditional customs, such as unsafe funerals/burials involving direct contact with body fluids of the dead due to the belief that the dead persons have to be well respected in order to be at peace with them [23,25].

### **Implication for COVID-19 transmission**

In view of the presence of pre-symptomatic and asymptomatic COVID-19 cases, the continued existence of these cultural and religious beliefs and practices outlined above has implications for COVID-19 transmission.

### **Poor perceived severity of the disease and low self-perceived risk**

Anecdotal evidence suggests that some Nigerians without confirmed COVID-19 still perceive the disease as “not severe” or “not real”. This perception is further fuelled by the number of persons confirmed as COVID-19 cases without symptoms [26]. These pre-symptomatic and asymptomatic COVID-19 cases do not believe that COVID-19 is real because they do not have symptoms. This has implication for COVID-19 transmission as the probability of adherence to COVID-19 public health preventive measures (such as good hand hygiene practices, good cough etiquette, wearing of face mask in public places, physical distancing and avoidance of crowded places) by persons who perceive COVID-19 as “not severe” or “not real” is very low [26].

Similarly, socio-cultural beliefs imbibe low self-perceived risk about COVID-19 in most people. This perception that they cannot be infected with COVID-19 increases their chances of exposure to the disease as they are also likely to ignore several public health preventive advisory and preventive measures outlined by the government and health officials.

In instances where individuals with poor perceived severity of COVID-19 or low self-perceived risk or both have suboptimal compliance to public health preventive advisory and preventive measures, the risk of been infected by unidentified or identified pre-symptomatic or asymptomatic COVID-19 cases is very high. This is particularly worse in crowded environments like social gatherings, markets, religious gatherings and so on, where intense community transmission of the disease can occur.

### **Resistance to treatment in accordance to case management protocol**

Several pre-symptomatic and asymptomatic confirmed cases resist isolation and treatment in the COVID-19 treatment facilities in several parts of Nigeria due to their belief and perception that only the sick, evidenced by manifestation of symptoms, are the ones who should be admitted in hospitals.



These cases do not see a need for the isolation because they believe they have no symptom or illness. Rather they may resort to self-medication or non-orthodox and traditional remedies, such as taking of herbs and visiting traditional healers, which have not been proven to be effective for the management of COVID-19, hence, leaving many of these cases as reservoir of the infection in the community.

Furthermore, resistance to treatment and case management could be heightened with the general information that the disease is novel and currently has no cure or vaccine for its prevention.

### Lack of trust in government, health agencies/institution and health expert opinions

With the current absence of specific treatment regimen for asymptomatic and pre-symptomatic confirmed cases of COVID-19 from the scientific world, many are returning to their beliefs in traditional institutions and religion to seek succor. The result is a lack of trust in the words and works of government, her agencies, representatives and experts as the source of solution to the pandemic. This is then manifested as disregard for government directives and outright deliberate disobedience to experts' advice and guides on the containment of COVID-19 [27].

Similarly, in some instances, people could refuse to report contacts of confirmed cases of COVID-19 to local health authorities and health facilities because they are seen as government institutions which they do not trust [27]. Literature has however reported contacts of confirmed cases, found during contact tracing, who tested positive to COVID-19 [28]. Therefore, it is obvious that many of these unreported contacts are just asymptomatic and pre-symptomatic cases waiting to be discovered. By not reporting them, they are left in the chain of transmission of COVID-19 within the community, thereby resulting in the spread of the disease.

### Recommendation

Because every region of Nigeria has its own multifaceted, culturally-specific context informing community perception, there is a need for the responses to be conducted in line with local realities, including culture and beliefs.

Hence, it is crucial that communities are engage in ways that reflects their shared beliefs and ideologies, and that approach should be reflected in planned behavioral change strategies. Furthermore, there is an urgent need to change the risk perception of majority of the populace on the COVID-19 disease. To do this, community engagement exercise needs to be strengthened. Opinion leaders such as influential religious leaders, traditional leaders and trade union leaders in whom most of the populace have more confidence, need to be brought on board to disseminate key message on COVID-19 preventive measures to the populace.

Also, public health authorities should substantially expand testing programs for COVID-19 to include persons who do not have symptom of COVID-19 and to isolate those who test positive using a strict enforcement protocol.

## References

1. Ibrahim RR, Oladipo HO (2020) Forecasting the spread of COVID-19 in Nigeria using Box-Jenkins Modeling Procedure. [Link: https://bit.ly/31oVuSM](https://bit.ly/31oVuSM)
2. World Health Organization (2020) Coronavirus disease (COVID-19) Situation Report-194. [Link: https://bit.ly/3gqztXY](https://bit.ly/3gqztXY)
3. NCDC (2020) COVID-19 situation report, situation report 150. [Link: https://bit.ly/2YsYlRh](https://bit.ly/2YsYlRh)
4. Paul A (2020) Tuberculosis and HIV responses threatened by COVID-19. *Lancet HIV* 5: e319- e320. [Link: https://bit.ly/2FTpvBN](https://bit.ly/2FTpvBN)
5. Ritu G, Sharma M (2020) 2019-nCoV pandemic: A disruptive and stressful atmosphere for Indian academic fraternity. *Brain Behav Immun* 88: 948-949. [Link: https://bit.ly/31oHrN1](https://bit.ly/31oHrN1)
6. Manik S, Sharma S (2020) The Rising Number of COVID-19 Cases Reflecting Growing Search Trend and Concern of People: A Google Trend Analysis of Eight Major Countries. *J Med Syst* 44: 117. [Link: https://bit.ly/3hqKgT4](https://bit.ly/3hqKgT4)
7. Nicola M (2020) The emotional impact of COVID- 19: from medical staff to common people. *Brain Behav Immun* 87: 23-24. [Link: https://bit.ly/3aRGD6a](https://bit.ly/3aRGD6a)
8. Adenomon MO, Maijamaa B, John DO (2020) On the Effects of COVID-19 outbreak on the Nigerian Stock Exchange performance: Evidence from GARCH Models. Preprints. [Link: https://bit.ly/2FPr43z](https://bit.ly/2FPr43z)
9. WHO. Coronavirus disease 2019 (COVID-19) Situation Report – 73. [Link: https://bit.ly/2FLCGVe](https://bit.ly/2FLCGVe)
10. Yu P, Zhu J, Zhang Z, Han Y (2020) A familial cluster of infection associated with the 2019 novel coronavirus indicating possible person-to-person transmission during the incubation period. *J Infect* 221: 1757-1751. [Link: https://bit.ly/3jaZexe](https://bit.ly/3jaZexe)
11. Huang R, Xia J, Chen Y, Shan C, Wu C (2020) A family cluster of SARS-CoV-2 infection involving 11 patients in Nanjing, China *Lancet Infect Dis* 20: 534-535. [Link: https://bit.ly/3ldTmVF](https://bit.ly/3ldTmVF)
12. Pan X, Chen D, Xia Y, Wu X, Li T, et al. (2020) Asymptomatic cases in a family cluster with SARS-CoV-2 infection. *Lancet Infect Dis* 20: 410-411. [Link: https://bit.ly/2EdUuYZ](https://bit.ly/2EdUuYZ)
13. Tong ZD, Tang A, Li KF, Li P, Wang HL, et al. (2020) Potential presymptomatic transmission of SARS-CoV-2, Zhejiang Province, China, 2020. *Emerg Infect Dis* 26. [Link: https://bit.ly/3gmYdQU](https://bit.ly/3gmYdQU)
14. Wei WE, Li Z, Chiew CJ, Yong SE (2020) Presymptomatic Transmission of SARS-CoV-2 -Singapore. *MMWR*.
15. Kimball A, Hatfield KM, Arons M, James A, Taylor J, et al. (2020) Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility – King County, Washington. *MMWR* 69: 377-381. [Link: https://bit.ly/3hkI3K9](https://bit.ly/3hkI3K9)
16. World Health Organization (2020) Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19) 16-24 February 2020 [Internet]. Geneva: World Health Organization. [Link: https://bit.ly/34sG9Cv](https://bit.ly/34sG9Cv)
17. Zhu C, Gao S, Xu Y, Yang XH, Ye FQ, et al. (2020) A COVID-19 case report from asymptomatic contact: implication for contact isolation and incubation management. *Infect Dis Poverty* 9: 70. [Link: https://bit.ly/3ja8Ubf](https://bit.ly/3ja8Ubf)
18. Oran DP, Topol EJ (2020) Prevalence of Aymptomatic SARS-COV-2 infection. *Annals of Internal Medicine*. [Link: https://bit.ly/34sTQSO](https://bit.ly/34sTQSO)
19. Wong G, Liu W, Liu Y, Zhou B, Bi Y, et al. (2015) MERS, SARS, and Ebola: The role of super-spreaders in infectious disease. *Cell Host & Microbes* 18: 398–401. [Link: https://bit.ly/3lhPOC2](https://bit.ly/3lhPOC2)
20. Bowale A, Abayomi A, Idris J, Omilabu S, Abdus-Salam I, et al. (2020)



Clinical presentation, case management and outcomes for the first 32 COVID-19 patients in Nigeria. Pan African Medical Journal 35: 24. [Link: https://bit.ly/32oLFUd](https://bit.ly/32oLFUd)

21. Olumba MO (2005) When Cultural Practices are Health Risks: The Dilemma of Female Genital Mutilation. Nigeria Journal of Health Education 13: 44-55.
22. Ojua TA, Ishor DG, Ndom PJ (2013) African Cultural Practices and Health Implications for Nigeria Rural Development. IRMBR 2: 176-183. [Link: https://bit.ly/32jWCGp](https://bit.ly/32jWCGp)
23. Adegoke AA (2008) Factors influencing health beliefs among people in the south West, Nigeria. Afr Research Rev 2: 177-197. [Link: https://bit.ly/3gwatyE](https://bit.ly/3gwatyE)
24. Champion VL, Skinner CS (2020) The health belief model 2008. In: Health behavior and health education theory, research, and practice [Internet]. San Francisco, CA: Jossey-Bass. 4<sup>th</sup> Edition. [Link: https://bit.ly/2EpVqsN](https://bit.ly/2EpVqsN)

25. Archibong EP, Enang EE, Bassey GE (2017) Witchcraft Beliefs In Diseases Causation And Health – Seeking Behaviour In Pregnancy Of Women In Calabar South – Nigeria. IOSR-JHSS 22: 24-28. [Link: https://bit.ly/3lll82P](https://bit.ly/3lll82P)

26. Evaborhene Aghogho Nelson. Coronavirus in Nigeria: A Case for Community Engagement. [Link: https://bit.ly/3lFk7ED](https://bit.ly/3lFk7ED)
27. Reuben RC, Danladi MMA, Saleh DA, Ejembi PE (2020) Knowledge, Attitudes and Practices towards COVID-19: An Epidemiological Survey in North-Central Nigeria. Journal of Community Health. [Link: https://bit.ly/2Qk2byX](https://bit.ly/2Qk2byX)
28. Daihai He, Zhao S, Lin Q, Zhuang Z, Cao P, et al. (2020) The relative transmissibility of asymptomatic COVID-19 infections among close contacts. Int'l J Infect Dis 145-147. [Link: https://bit.ly/3hoEqSt](https://bit.ly/3hoEqSt)

### Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

#### Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROMEO, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

**Submit your articles and experience a new surge in publication services**  
(<https://www.peertechz.com/submission>).

*Peertechz journals wishes everlasting success in your every endeavours.*

**Copyright:** © 2020 Isere EE, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Isere EE, Adejugbagbe AM, Oladoyin V, Abiona S, Omorogbe EN (2020) Pre-symptomatic and asymptomatic COVID-19 cases in Nigeria amidst prevailing socio-cultural beliefs and practices: Implication for COVID-19 transmission and way forward. Arch Prev Med 5(1): 039-042.  
DOI: <https://dx.doi.org/10.17352/apm.000018>