

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:IJDCR-2-117

[A Japanese Case of Nonpustular Annular Psoriasis](#)

Published On: September 24, 2016 | Pages: 019 - 020

Author(s): Hidetoshi Takahashi*

Introduction Nonpustular annular psoriasis is a rare form of psoriasis with clinically annular ring-shaped configuration [1]. In contrast to recurrent circinate erythematous psoriasis (Psoriasis a type d'erytheme circine recidivant de Bloch) [2], which is the mildest form of pustular psoriasis, no evidence of histopathological Kogoi's spongiform pustule formation i ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000017

[Open Access](#) [Research Article](#) PTZAID:IJDCR-2-113

[Metabolic Associations with Skin Tags](#)

Published On: February 20, 2016 | Pages: 003 - 011

Author(s): Azar Hadi Maluki*, Aala Abidmuslim Abdullah

Introduction Part 1 / skin tags Synonyms: Soft warts, achrochordon[1], cutaneous tag, papilloma colli, fibroma pendulum, cutaneous papilloma, fibroma molluscum, templeton skin tags [2], fibroepithelial polyps, pedunculated soft fibromas, filiform soft fibromas [3]. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000013

Case Report

[Open Access](#) [Case Report](#) PTZAID:IJDCR-2-115

[Systemic Sclerosis Associated Angiosarcoma: A Case Report and Review of the Literature](#)

Published On: August 27, 2016 | Pages: 014 - 017

Author(s): Brittany Carter*, Christine Jaworsky, Monte Fox

Introduction Cutaneous angiosarcoma is an uncommon malignant vascular tumor with a highly variable presentation. It has a predilection for elderly Caucasian men, with 50% of tumors occurring primarily on the head and neck [1].

Cutaneous angiosarcomas may also occur in sites with chronic lymphedema, areas of prior radiation therapy, and in association with vascular ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000015

Letter to Editor

[Open Access](#) | [Letter to Editor](#) | PTZAID:IJDCR-2-116

[Localized Myxedema on the Upper Eyelids in a Patient of Hypothyroidism](#)

Published On: September 23, 2016 | Pages: 018 - 018

Author(s): Hidetoshi Takahashi*, Akemi Ishida-Yamamoto and Hajime Iizuka

Letter Localized myxedema may be observed in hypo/hyperthyroidism [1]. However, the affected sites are usually pretibial area and the lesion on the eye lid is rare. We report a case of myxedema on the bilateral upper eyelids, which was a diagnostic clue of hypothyroidism associated with Hashimoto disease. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000016

[LCDC \(Leprosy Cases Detection Campaign\) - Aiming at Cent Percent Detection of Leprosy Cases in the Community Based on a Line of Pulse Polio Campaign in Leprosy Endemic State](#)

Published On: May 02, 2016 | Pages: 012 - 013

Author(s): Sunil Vilasrao Gitte*

Letter Sir, India alone, contributes about 58.8% and 81.2% of the newly detected cases of leprosy, globally and in the South East Asian region, respectively [1]. Cases were detected mainly by passive surveillance. Detection of leprosy cases will lead to a diminution of the source of infection in the community, mainly because undetected and untreated cases will t ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/2455-8605.000014](#)

[Nail Fold Capillaroscopic Findings in Elderly Patient with Acrocyanosis](#)

Published On: January 08, 2016 | Pages: 001 - 002

Author(s): Maria Bibiana Leroux*

Dear Editor The Raynaud´s phenomenon (RP) is well known as the first sign of systemic scleroderma (SS) and related disorders. Its early diagnosis leads to a better understanding of the disease and favorable prognosis. ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/2455-8605.000012](#)