

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:IJDCR-3-124

[A Study on clinical patterns of mucocutaneous candidiasis in immunosuppressed patients](#)

Published On: December 12, 2017 | Pages: 032 - 034

Author(s): Subhashini Mohan*

Candidiasis is the most common superficial fungal infection in immunocompromised patients. With the advent of wide spectrum of immunosuppressive drug and increase in HIV patients, there is a change in the epidemiology and clinical presentation of mucocutaneous candidiasis. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000024

[Open Access](#) [Research Article](#) PTZAID:IJDCR-3-123

[Role of Helicobacter Pylori in Chronic Urticaria among Egyptian Patients with Dyspepsia: A case-control study](#)

Published On: October 31, 2017 | Pages: 026 - 031

Author(s): Al-Sadat Mosbeh*

Chronic urticaria is one of the most frequent skin diseases and still its etiology is recognized only in a minority of cases. Some recent studies point out to infections due to Helicobacter Pylori as being of major importance in the pathogenesis of chronic urticaria. This study aimed to find out the association of chronic urticaria with H. pylori ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000023

[Open Access](#) [Research Article](#) PTZAID:IJDCR-3-122

[Profile of Bacterial Dermohypodermatitis in Senegal over a 30-year period](#)

Published On: September 21, 2017 | Pages: 022 - 025

Author(s): Diallo M*, Niasse M, Diatta BA, Diop A, Ndiaye M, Ly F and Dieng MT

Introduction: The increase of both the frequency and severity of the bacterial dermohypodermatitis (BDH) in Senegal suggests the existence of other risk factors outside of those classically known to be involved. Our study aimed to clarify the epidemiological, clinical and risk factors of BDH in Senegal. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000022

[Open Access](#) | [Research Article](#) | PTZAID:IJDCR-3-118

[Skin Disorders in Adult Obese Egyptian Patients: A Comparative Study](#)

Published On: April 25, 2017 | Pages: 001 - 007

Author(s): Ensaf Mohamed Abdel Maguid, Sara Mohamed Awad*, Maha Mohammed Saad

Background: Obesity is a major public health concern worldwide. Limited number of studies on the relationship between obesity and skin disease are available ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000018

Review Article

[Open Access](#) | [Review Article](#) | PTZAID:IJDCR-3-121

[Comparison of safety and efficacy of Oral Terbinafine with Amphotericin B gel and Sertaconazole cream for the treatment of Tinea Corporis and its effect on quality of life of patients](#)

Published On: September 09, 2017 | Pages: 018 - 021

Author(s): Sumedha Sirohi, Sumit Bhutani, Cherry Cheema, Gurleen Kaur, S P Singh and Prithpal S Matreja*

Tinea Corporis is a superficial fungal infection affecting 20-25% of the world's population. Management involves using topical and oral antifungal drugs. Terbinafine has been used commonly for the treatment of tinea corporis to prevent the emergence of resistance combination of drugs is used. Hence, this study was designed to compare the safety and efficacy of Sert ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000021

[Open Access](#) | [Review Article](#) | PTZAID:IJDCR-3-120

[Topical Curcumin: A Review of Mechanisms and uses in Dermatology](#)

Published On: July 06, 2017 | Pages: 010 - 017

Author(s): Heng MCY

Curcumin, the active ingredient in the spice turmeric, has been used in many Eastern countries for its known anti-inflammatory activity. Recently, ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000020

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:IJDCR-3-119

[A Case of Jessner's Lymphocytic Infiltrate and Cutaneous B-Cell Follicular Lymphoma: Coexistence or Transformation?](#)

Published On: June 15, 2017 | Pages: 008 - 009

Author(s): Bryan Murphy*, Helen Hunter and Jennifer Somerville

Jessner's lymphocytic infiltrate of the skin is a rare, benign cutaneous disorder. Difficulty arises in differentiating it from cutaneous lymphoma on histology, thus requiring immunophenotyping, however the literature lacks reports demonstrating malignant transformation or dual pathology. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8605.000019